

Supplementary Table 2 (A). Relationship between stress urinary incontinence and self-administered questionnaires 6 months after surgery

Variable	Satisfaction with treatment question					Total	
	Very satisfied	Rather satisfied	Neutral	Rather unsatisfied	Very unsatisfied		
Patients without SUI	773 (73.4)	218 (20.7)	33 (3.1)	25 (2.4)	4 (0.4)	1,053 (100)	
Patients with SUI	7 (53.8)	5 (38.5)	0 (0)	1 (7.7)	0 (0)	13 (100)	
P-value	0.336						
Variable	Overall response assessment			Total			
	Improved	No change	Aggravated				
Patients without SUI	1,044 (99.1)	5 (0.5)	4 (0.4)	1,053 (100)			
Patients with SUI	12 (92.3)	0 (0)	1 (7.7)	13 (100)			
P-value	0.001*						
Variable	Willingness to undergo surgery question					Total	
	Definitely	Most likely	Somewhat likely	Not likely	Mostly not likely		Never
Patients without SUI	699 (66.4)	196 (18.6)	107 (10.1)	29 (2.8)	15 (1.4)	7 (0.7)	1,053 (100)
Patients with SUI - No. ()	5 (38.5)	5 (38.5)	2 (15.4)	1 (7.7)	0 (0)	0 (0)	13 (100)
P-value	0.318						

Values are presented as number (%).

SUI, stress urinary incontinence.

P-value, difference between stress urinary continence and self-administered questionnaires 6 months after surgery.

*P < 0.05, statistically significant differences.

Supplementary Table 2 (B). Relationship between urgency urinary incontinence and self-administered questionnaires 6 months after surgery

Variable	Satisfaction with treatment question					Total	
	Very satisfied	Rather satisfied	Neutral	Rather unsatisfied	Very unsatisfied		
Patients without UUI	775 (73.5)	218 (20.7)	33 (3.1)	24 (2.3)	4 (0.4)	1,054 (100)	
Patients with UUI	5 (41.7)	5 (41.7)	0 (0)	2 (16.7)	0 (0)	12 (100)	
P-value	0.006*						
Variable	Overall response assessment			Total			
	Improved	No change	Aggravated				
Patients without UUI	1,044 (99.1)	5 (0.5)	5 (0.5)	1,054 (100)			
Patients with UUI	12 (100)	0 (0)	0 (0)	12 (100)			
P-value	0.944						
Variable	Willingness to undergo surgery question					Total	
	Definitely	Most likely	Somewhat likely	Not likely	Mostly not likely		Never
Patients without UUI	699 (66.3)	199 (18.9)	107 (10.2)	29 (2.8)	13 (1.2)	7 (0.6)	1,054 (100)
Patients with UUI	6 (50.0)	3 (25.0)	1 (8.3)	1 (8.3)	1 (8.3)	0 (0)	12 (100)
P-value	0.279						

Values are presented as number (%).

UUI, urgency urinary incontinence.

P-value, difference between stress urinary continence and self-administered questionnaires 6 months after surgery.

*P < 0.05, statistically significant differences.